

Measuring the Psychological Impacts of Prison-Based Dog Training Programs and In-Prison Outcomes for Inmates

The Prison Journal

1–16

© 2019 SAGE Publications

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0032885519894657

journals.sagepub.com/home/tpj

Erin Flynn¹, Katie Massey Combs²,
Jaci Gandenberger¹, Philip Tedeschi¹,
and Kevin N. Morris¹

Abstract

As interest grows in programs that improve prison inmates' behavior and psychosocial well-being, any such interventions must be rigorously examined and their underlying mechanisms for change must be understood. This pilot study examined the use of prison-based dog training programs across Washington State Department of Corrections facilities for their impacts on inmates' infraction rates. The study also compared levels of empathy, self-efficacy, and anxiety between program participants and nonparticipants. Findings indicated that prison dog program participants' infraction rates improved and that participants had lower levels of anxiety than nonparticipants.

Keywords

prison, prison-based animal programs, prison dog programs, infraction rates

¹University of Denver, CO, USA

²University of Colorado Boulder, USA

Corresponding Author:

Kevin N. Morris, Institute for Human-Animal Connection, Graduate School of Social Work,
University of Denver, 2148 High Street, Denver, CO 80208, USA.

Email: kevin.morris@du.edu

Introduction

An estimated 2.3 million people were incarcerated in the United States in 2016, a significantly higher number than that of any other country in the world (Kaeble & Cowhig, 2018; Warren et al., 2008). Although efforts are underway to reduce prison populations, it is essential that any such efforts maintain safety for staff, inmates, and the wider population (Gelb & Stevenson, 2017; Grawert et al., 2017). Programs that reduce inmate in-prison infraction rates, particularly those that also improve inmates' psychosocial well-being, can be an effective way to maintain staff and inmate safety within correctional facilities while also reducing the likelihood of recidivism upon release (Cochran et al., 2014; Fournier et al., 2007; Steiner & Meade, 2016). Approximately 95% of prisoners are eventually released from prison, so successfully addressing inmate infraction and psychosocial well-being is an important component in improving the safety of both correctional facilities and broader society (Hughes & Wilson, 2018).

Policymakers have explored a variety of approaches to achieve these goals, including more conventional programs such as mental health therapy and addictions treatment as well as alternative programs ranging from meditation retreats to Shakespearean theater (Council of State Governments Justice Center, 2014; Heard et al., 2013; Perelman et al., 2012). One promising but understudied approach is the implementation of prison-based dog programs (PDPs), in which inmates become dog handlers who train the animals for adoption or service work (Van Wormer et al., 2017). Current research indicates that these programs improve inmates' well-being and reduce both infraction rates and recidivism (Cooke & Farrington, 2016; Van Wormer et al., 2017). However, that research has been limited by a number of factors, including small sample sizes, single-site studies, the use of nonvalidated measures, heavy reliance on qualitative data and anecdotal reports, omission of details regarding research processes that makes replication difficult, and the complexity of performing randomized controlled trials in prison settings (Cooke & Farrington, 2014b, 2016; Fournier et al., 2007; Mulcahy & McLaughlin, 2013; Van Wormer et al., 2017).

One recent study addressed some of these concerns using a larger sample size ($n = 1,001$) to compare infraction rates of participants in PDPs and a group of nonparticipating inmates statistically matched across several criteria, including average incarceration length, custody level, and baseline levels of outcome variables (Van Wormer et al., 2017). This study, the largest and one of the few to use a statistically matched comparison group to date, found that inmates who participated in PDPs had significantly lower serious and violent infraction rates than matched inmates who did not participate in them.

These findings support the use of such programs to create “safer and healthier” prison environments by reducing infraction rates (Van Wormer et al., 2017, p. 537). However, the underlying psychosocial mechanisms affecting the improved infraction rates are still poorly understood and warrant further study so that the impacts of PDPs can be optimized.

Literature Review

Although the incorporation of animals in prisons and other institutional settings is frequently considered innovative, it is not new. The first documented use of animal-assisted therapy (AAT) occurred in 1792, in which farm animals were used to teach mental patients self-control (Furst, 2011). Animal programs were introduced into prisons in 1981, starting at the Washington State Corrections Center for Women, and now exist in all 50 states in the United States as well as several foreign countries (Cooke & Farrington, 2014b; Strimple, 2003). Today, prison-based animal programs (PAPs) vary widely in scope, aim, eligibility, and even the species of animals involved. Although this variety poses challenges to studying the efficacy of PAPs, it is likely a contributing factor to their popularity (Cooke & Farrington, 2014b).

The Washington State Department of Corrections (WADOC) continues to be a leader in implementing these programs. Every WADOC facility has at least one animal training or adoption program, which are intended to provide dog training, care, and adoption services to local communities while improving inmates’ responsibility and lowering their rates of violence and recidivism (LeRoy et al., 2012; WADOC, 2016).

WADOC’s dog programs are not designed to be therapeutic in nature, but rather to teach inmates skills in dog training, grooming, and general care (Van Wormer et al., 2017). However, findings from exploratory and qualitative studies suggest that PDPs may yield positive therapeutic impacts, such as increased self-efficacy, empathy, and emotional well-being (Cooke & Farrington, 2014b; Minton et al., 2015; Richardson-Taylor & Blanchette, 2001). Although intriguing, the current research on the potential therapeutic benefits of PDPs—like the research on PAPs more generally—has multiple limitations, as described above, particularly single-site studies and the use of nonvalidated measures. This pilot study aims to partially address current gaps by applying widely used, highly validated measures of empathy, self-efficacy, and anxiety to compare PDP participants and nonparticipants in 10 WADOC prisons. The three psychosocial variables examined in this study were chosen due to their inclusion in previous studies of participants in PDPs and association with inmates’ infraction rates and rehabilitation.

Programs that aim to increase empathy, defined here as “the ability to understand and share the feelings of another” (Yogev, 2012, p. 61) among inmates are common in many prisons because this psychosocial parameter is a central component of prosocial behavior and is inversely related to violence, criminal offenses, and recidivism (Bock & Hosser, 2014; Hepper et al., 2014; Jolliffe & Farrington, 2004; Miller & Eisenberg, 1988; Van Langen et al., 2014). Although it remains unclear whether interactions with an animal increase empathy, human bonds with animals are positively correlated with empathy (Beetz et al., 2012). Furthermore, increased empathy toward dogs and other humans seems to occur among participants in PDPs and is a perceived benefit found in many surveys of both inmates and prison staff (Cooke & Farrington, 2014b, 2016; Furst, 2011; Minton et al., 2015).

Self-efficacy, or the “belief in one’s capabilities to mobilize the motivation, cognitive resources and courses of action needed to meet given situational demands” (Gist & Mitchell, 1992, p. 184), is frequently noted as a benefit of PDPs in qualitative interviews (Cooke & Farrington, 2016; Furst, 2011; Lai, 1998; Richardson-Taylor & Blanchette, 2001; Strimple, 2003). For example, in an exploratory study conducted by Cooke and Farrington (2014b), five of the 11 PDP staff stated that self-efficacy was improved by participating in these programs, despite that their questionnaire did not directly ask about self-efficacy. Similar findings emerged from interviews with inmates who participated in PDPs (Cooke & Farrington, 2014a; Leonardi et al., 2017). General self-efficacy gained through prison programs is a powerful behavior change determinant because an individual’s expectations of their own ability inform their decision to perform a behavior, expend effort, and persevere in the face of adverse conditions (Bandura, 1977, 1982). If PDPs are particularly effective in increasing inmate self-efficacy, it could have significant implications for inmate rehabilitation efforts.

High levels of anxiety or “tension, apprehension, nervousness, and worry,” accompanied by “arousal of the autonomic nervous system” (Spielberger, 1983, p. 4), among inmates can lead to increased conflicts with correctional officers and other inmates (MacKenzie, 1987). Contact with animals correlates positively with improved human health and reduced stress, which tend to influence levels of anxiety (Beetz et al., 2012, Grinde & Patil, 2009; Wilson, 1984). PDPs have been frequently reported as increasing happiness and reducing stress among program participants as well as reducing the overall stressfulness of prison environments (Cooke & Farrington, 2014a, 2016; Furst, 2011; Harkrader et al., 2004; Richardson-Taylor & Blanchette, 2001; Strimple, 2003). In one survey, 76% ($n = 23$) of participants reported a clear reduction in stress, although others noted that their stress was sometimes increased due to having to care for dogs in lockdown situations (Minton

et al., 2015). Furthermore, anecdotal reports have linked PDPs to improvements in individual participants' anxiety symptoms as well as other potential anxiety markers such as high blood pressure (Osborne & Blair, 2003).

Further research is needed regarding the psychosocial impacts of PDPs (Cooke & Farrington, 2016; Van Wormer et al., 2017). The aim of this pilot study was to further examine the impact of PDPs on infraction rates as well as psychosocial outcomes, specifically empathy, self-efficacy, and anxiety, and to explore whether validated self-report instruments can be used to examine the efficacy of dog programs within a prison setting. If so, this supports further research to implement these tools on a broader scale, allowing for a more consistent and reliable set of data to be used to analyze the impacts of these programs over time.

Method

Participant Selection

This pilot study included 229 male and female inmates from 10 of 12 WADOC facilities across all security levels. To be eligible to participate, inmates were required to be at least 18 years of age and proficient in reading English. The recruitment and consent process consisted of two phases. First, 1 month prior to data collection, approximately 450 inmates were identified by the research team as potential participants. Each inmate was sent a recruitment flyer by WADOC staff; individuals were able to ask the staff questions prior to the researcher being on site. The second phase took place on the day of the survey session. Inmates eligible to participate responded to a voluntary call out to the survey session. Respondents were gathered in groups and guided through a recruitment and consent script and allowed to ask questions regarding their potential participation. Those who agreed to participate and signed the consent form were included in the study. Participants were tracked by their WADOC identification numbers. Prior to data analysis, these numbers were replaced with randomly assigned study identification numbers.

Of the total of 229 inmates in the study, 150 were in the PDP group and 79 were included as comparison nonparticipants. For the sake of the study, inmates designated as participants were those who were ever in a PDP; they may not have been actively participating at the time of the survey data collection. Descriptive statistics of the sample by intervention and comparison groups were collected (Table 1). Of the participants, 85.6% identified as White, 9.6% as Black, and 4.4% as Other (Asian and American Indian.) One of the 10 prisons studied was a women's facility, with women making up 6.60% of the total participants. Inmates' risk levels were determined using

Table 1. Descriptions of Participant Demographics.

	Total (229), %/M (SD)	Intervention (n = 150), %/M (SD)	Comparison (n = 79), %/M (SD)	Chi- square/U test
Race				
White	85.6%	86.0%	84.8%	4.99
Black	9.6%	10.0%	8.9%	
Other (Asian, American Indian)	4.4%	4.0%	6.3%	
Gender—Female	6.6%	9.3%	1.3%	5.50*
Age at survey data collection	39.9 (12.3)	39.4 (13.0)	40.9 (11.0)	
Risk level				
Low	33.60%	37.30%	26.60%	2.79
Moderate	14.40%	13.30%	16.50%	
High	51.90%	49.40%	57.00%	
Pre-PDP infraction rate (<i>Mdn</i>)	0.68	0.54	0.68	3,939.00
Median post-/during-PDP infraction rate (<i>Mdn</i>)	0.59	0.34	1.01	4,155**

Note. PDP = prison-based dog program.

* $p < .05$. ** $p < .01$.

standard WADOC protocol, which calculates scores based on a variety of weighted offender risk factors and classified inmates based on the score achieved (Washington State Institute for Public Policy, 2008). Of the total participants, 33.6% were rated as low risk, 14.4% as medium, and 51.9% as high. The categories of “high violent,” “high property,” and “high drug” risk levels were collapsed into “high.” The infraction rates listed represent the number of infractions divided by the number of days in prison. No significant differences were found between the intervention and comparison groups in terms of race, risk level, or preprogram infraction rates.

Measures

Infractions

Change in infraction rates before and during the PDP was assessed using administrative data from WADOC. These data included infractions and entry and exits for each participating inmate from June 2011 to June 2017. Infraction rates were calculated, with the total number of infractions divided

Table 2. Wilcoxon Signed-Ranks Tests on Pre- and Postinfracton Rates.

	Preprogram: Median infracton rate	Post-/during- program: Median infracton rate	z-test	Significance
Intervention	0.68	0.34	-2.52	.012
Comparison	0.54	1.01	-1.47	.141

by the total number of days in prison during the two periods of the study (pre-PDP and during-PDP). Because participants had varying start dates in the PDP and may have exited and reentered prison between June 2011 and June 2017, the number of days in the program was calculated based on individual PDP start dates and actual days in prison. For the comparison group, since the first implementation of the PDP was June 22, 2014, with 25% of the intervention group starting on this date, June 22, 2014, was used as the cutoff between pre-PDP and “during” PDP. The mean number of infractions pre-PDP was 1.55 ($SD = 2.14$) and during the PDP was 1.60 ($SD = 2.66$). The infractions rates were not normally distributed, thus medians are reported, and nonparametric tests were utilized for infracton rate analyses. The comparative pre- and postinfracton rates between the dog and no dog groups were compared (Table 2).

Empathy

The Brief Interpersonal Reactivity Index (B-IRI) was used to measure empathy level (Ingoglia et al., 2016). The full-length Interpersonal Reactivity Index (IRI) consists of 28 Likert-type scale questions covering four subscales—Fantasy, Empathic Concern, Perspective Taking, and Personal Distress—that measure both cognitive and affective empathy (Davis, 1980, 1983). It is the most widely used self-report measure of empathy over the last 20 years and has been employed in prison populations (Gerdes et al., 2010; Jolliffe & Farrington, 2004; Loinaz et al., 2018). However, several studies have indicated that inmates, particularly those with lower IQs and weaker verbal skills, may have difficulty with some of the questions to an extent that lowers the IRI’s reliability with those populations (Beven et al., 2004; Lauterbach & Hosser, 2007). To address those concerns, Ingoglia et al. (2016) developed the 16-question B-IRI as a shorter, more accessible version for special populations with low literacy skills. A series of studies across three independent samples demonstrated that the B-IRI preserves “a reasonable scale reliability and validity” compared with the full-length IRI (Ingoglia

et al., 2016, p. 9). Strong reliability was indicated for Empathic Concern ($\alpha = .757$), Perspective Taking ($\alpha = .798$), and Fantasy ($\alpha = .760$). Reliability was not as strong on the Personal Distress subscale ($\alpha = .570$), thus it was not included in the analysis.

Self-Efficacy

The General Self-Efficacy Scale (GSES) was used to measure self-efficacy (Schwarzer & Jerusalem, 1995). It contains 10 statements, written to a basic literacy level that the respondent scores on a Likert-type scale from “not true” to “completely true,” and is designed to broadly assess a person’s belief in their ability to cope with daily challenges and to adapt after experiencing common stressful events. The GSES is designed for a broad adult population and has been validated and implemented in a variety of environments, including prison education programs (Allred et al., 2013; Luszczynska et al., 2005; Luszczynska et al., 2010; Schwarzer & Jerusalem, 1995). The GSES had strong reliability within this sample ($\alpha = .884$).

Anxiety

The State/Trait Anxiety Index for Adults (STAI-AD) was used to measure anxiety (Spielberger, 1972, 1983; Spielberger et al., 1970). The STAI-AD, which is written to a sixth grade reading level, has been validated for measuring temporary, or “State,” versus inherent, or “Trait,” anxiety levels in individuals. It has been successfully employed in a wide variety of environments, including prisons (Barnes et al., 2002; Chen et al., 2016; Kabacoff et al., 1997). The State items ($\alpha = .942$) and the Trait items ($\alpha = .920$) had very strong reliability.

Participants completed the B-IRI, GSES, and STAI-AD at a single point in time, 3 years after the start of the study period (Figure 1).

Analysis

The change in infraction rates from pre-PDP to during-PDP was also assessed. Because the rate of infractions was not normally distributed, nonparametric tests were employed. The Mann–Whitney U test for nonparametric variables was used to examine differences pre-PDP between the infraction rates between the two groups, and the Wilcoxon rank-sum test to examine change in infraction rates from pre-PDP to during-PDP for each group. Independent sample t -tests were conducted to examine differences in psychosocial measures between the intervention and comparison groups. Differences in




Figure 1. Infraction rate analysis timeline.

infraction rates were examined between the intervention and comparison groups pre-PDP.

Results

Infraction Rates

A Mann–Whitney U test indicated that infraction rates prior to anyone experiencing the PDP were similar for the intervention and the comparison groups, $U = 3,939.00, p = .577$, although the intervention group had significantly lower infraction rates post-/during-PDP, $U = 4,155.00, p < .001$ (Table 2). Wilcoxon signed-ranks tests indicated that intervention participants had a significant decrease in infraction rates from pre-PDP to post-/during-PDP, whereas inmates not in the dog program experienced no significant change during these two periods (Table 2).

Psychosocial Measures

Inmates in the two groups had similar scores in all but one of the B-IRI subscales; however, PDP participants had higher scores on the empathy subscale

Table 3. Independent Samples *t*-tests on Psychosocial Measures.

	Intervention			Comparison			<i>t</i> -test	Cohen's <i>d</i>
	<i>n</i>	<i>M</i>	<i>SD</i>	<i>n</i>	<i>M</i>	<i>SD</i>		
GSES	149	3.23	0.47	78	3.15	0.55	1.16	0.156
S_Mean	149	1.54	0.48	78	1.70	0.61	-2.09*	0.292
T_Mean	149	1.91	0.49	78	1.98	0.56	-0.89	0.133
B-IRI Empathy	150	4.00	0.75	78	3.81	0.88	1.73***	0.232
B-IRI Perspective Taking	150	3.42	0.94	78	3.42	0.93	0.01	0
B-IRI Fantasy	150	3.09	1.00	78	3.09	0.96	0.04	0

Note. GSES = General Self-Efficacy Scale; B-IRI = Brief Interpersonal Reactivity Index.

* $p < .05$. *** $.5 > p < .1$.

of the B-IRI that approached significance ($p = .09$). No differences were found between the two groups on the GSES scores or the Trait anxiety subscale, but PDP participants reported statistically significant lower scores on the State subscale ($p = .03$) (Table 3).

Discussion

This study replicated previous findings across a different time frame indicating that participation in PDPs lowers infraction rates among participants compared with inmates who are not in the programs. The similar infraction rates between the two groups prior to entering PDPs and the maintenance of these rates in the non-PDP group across the entire study period indicate that programs involving dogs decrease problem behaviors. In this regard, it is important to note that infractions often lead to dismissal from PDPs. The study also demonstrated that psychosocial data that might further explain the mechanisms of change for PDPs can be collected in prison environments using validated instruments. No statistically significant difference was measured in self-efficacy between the participants and nonparticipants in PDPs, but the dog program inmates had lower anxiety ($p = .03$) and were approaching statistical significance for higher empathy ($p = .09$). The effect sizes for both differences were small. Because these data were only measured at a single time point, further research is necessary to state whether PDP participation lowers anxiety or if inmates with lower levels of anxiety are more likely to participate in PDPs. However, the fact that there was no difference in the Trait anxiety scores between the two groups suggests that PDPs may be effective in lowering anxiety for participants.

Limitations

This study has several strengths, including the inclusion of both male and female inmates from multiple prisons across security levels, as well as the use of well-validated psychosocial instruments. Although the participants did not differ on most major demographic variables (see Table 1), other unobserved differences likely existed related to risk, motivation, or other factors. Furthermore, the single time point does not allow determination of whether inmates in PDPs experience changes in psychosocial health or if individuals with these characteristics are attracted to such programs. However, the pilot findings support the expansion of these types of studies to better understand the impacts of PDPs on inmate outcomes.

Because all inmates are typically exposed to dogs within prisons that have PDPs, it is possible that some of the impacts of these programs were obscured: the mere presence of dogs may affect all inmates being studied, including those in the comparison group. The data may be further obscured because those in the intervention group did not necessarily have a dog at the time of data collection. This further bolsters the value of future research that uses these or similar well-validated measures to examine psychosocial data across multiple time points—to gather and assess data on both dog handlers and nonhandlers exposed to PDPs.

An additional point of note is that, while many of the inmates were eager to discuss their experiences with the PDPs, participants frequently expressed disappointment or frustration when the research consisted of psychosocial assessments rather than more qualitative opportunities to speak directly about the programs. Although further high-quality quantitative research is nonetheless called for, it would ideally be paired with qualitative approaches using mixed methods designs that also capture a nuanced understanding of the inmates' perceptions of PDPs.

Summary and Future Directions

The demonstration that potential psychosocial change mechanisms can be identified using validated instruments for inmates participating in PDPs supports further research aimed at delineating those mechanisms. By developing a better understanding of how PDPs impact participants, those leading or developing these programs can better optimize their impacts on inmate outcomes. Although these programs are increasingly common across the United States and internationally, little work has been done to establish best practices for the inmates, staff, or dogs involved. Such future work should include the further use of validated psychosocial measures, ideally across a greater number of sites and over multiple time points.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research and/or authorship of this article: The authors received financial support for the research, authorship, and publication of this article through a University of Denver Professional Research Opportunities for Faculty grant (DU Award 89959).

References

- Allred, S. L., Harrison, L. D., & O'Connell, D. J. (2013). Self-efficacy: An important aspect of prison-based learning. *The Prison Journal, 93*(2), 211–233. <https://doi.org/10.1177/0032885512472964>
- Bandura, A. (1977). Self-efficacy: Toward a unifying theory of behavioral change. *Psychological Review, 84*(2), 191–215.
- Bandura, A. (1982). Self-efficacy mechanism in human agency. *American Psychologist, 37*, 122–147.
- Barnes, L. B., Harp, D., & Jung, W. S. (2002). Reliability generalization of scores on the Spielberger State-Trait Anxiety Inventory. *Educational and Psychological Measurements, 62*(4), 603–618. <https://doi.org/10.1177/0013164402062004005>
- Beez, A., Uvnas-Moberg, K., Julius, H., & Kortschal, L. (2012). Psychosocial and psychophysiological effects of human-animal interactions: The possible role of oxytocin. *Frontiers in Psychology, 3*, Article 234. <https://doi.org/10.3389/fpsyg.2012.00234>
- Beven, J. P., O'Brien-Malone, A. O., & Hall, G. (2004). Using the Interpersonal Reactivity Index to assess empathy in violent offenders. *International Journal of Forensic Psychology, 1*(2), 33–41.
- Bock, E. M., & Hosser, D. (2014). Empathy as a predictor of recidivism among young adult offenders. *Psychology, Crime & Law, 20*(2), 101–115. <https://doi.org/10.1080/1068316X.2012.749472>
- Chen, X. J., Hannibal, N., & Gold, C. (2016). Randomized trial of group music therapy with Chinese prisoners: Impact on anxiety, depression, and self-esteem. *International Journal of Offender Therapy and Comparative Criminology, 60*(9), 1064–1081. <https://doi.org/10.1177/0306624X15572795>
- Cochran, J. C., Mears, D. P., Bales, W. D., & Stewart, E. A. (2014). Does inmate behavior affect post-release offending? Investigating the misconduct-recidivism relationship among youth and adults. *Justice Quarterly, 31*(6), 1044–1073. <https://doi.org/10.1080/07418825.2012.736526>
- Cooke, B. J., & Farrington, D. P. (2014a). The effects of dog-training programs: Experiences of incarcerated females. *Women & Criminal Justice, 25*(3), 201–214. <https://doi.org/10.1080/08974454.2014.909763>

- Cooke, B. J., & Farrington, D. P. (2014b). Perceived effects of dog-training programmes in correctional settings. *The Journal of Forensic Practice, 16*(3), 171–183. <https://doi.org/10.1108/JFP-08-2013-0041>
- Cooke, B. J., & Farrington, D. P. (2016). The effectiveness of dog-training programs in prison: A systematic review and meta-analysis of the literature. *The Prison Journal, 96*(6), 854–876. <https://doi.org/10.1177/0032885516671919>
- Council of State Governments Justice Center. (2014). *Reducing recidivism: States deliver results*. [https://csgjusticecenter.org/wp-content/uploads/2014/06/Reducing Recidivism_StatesDeliverResults.pdf](https://csgjusticecenter.org/wp-content/uploads/2014/06/Reducing_Recidivism_StatesDeliverResults.pdf)
- Davis, M. H. (1980). A multidimensional approach to individual differences in empathy. *JSAS Catalog of Selected Documents in Psychology, 10*, 85.
- Davis, M. H. (1983). Measuring individual differences in empathy: Evidence for a multidimensional approach. *Journal of Personality and Social Psychology, 44*(1), 113–126. <https://doi.org/10.1037/0022-3514.44.1.113>
- Fournier, A. K., Geller, E. S., & Fortney, E. E. (2007). Human-animal interaction in a prison setting: Impact on criminal behavior, treatment progress, and social skills. *Behavior and Social Issues, 16*(1), 89–105.
- Furst, G. (2011). *Animal programs in prison: A comprehensive assessment*. Lynne Rienner Publishers.
- Gelb, A., & Stevenson, P. (2017). *U.S. adult incarceration rate declines 13% in 8 years*. The Pew Charitable Trusts. <http://www.pewtrusts.org/en/research-and-analysis/analysis/2017/01/12/us-adult-incarceration-rate-declines-13-percent-in-8-years>
- Gerdes, K. E., Segal, E. A., & Lietz, C. A. (2010). Conceptualizing and measuring empathy. *British Journal of Social Work, 40*(7), 2326–2343. <https://doi.org/10.1093/bjsw/bcq048>
- Gist, M., & Mitchell, T. R. (1992). Self-efficacy: A theoretical analysis of its determinants and malleability. *Academy of Management Review, 17*(2), 183–211. <https://doi.org/10.5465/AMR.1992.4279530>
- Grawert, A., Camhi, N., & Chettiar, I. M. (2017). *A federal agenda to reduce mass incarceration*. Brennan Center for Justice. <https://www.brennancenter.org/publication/federal-agenda-reduce-mass-incarceration>
- Grinde, B., & Patil, G. G. (2009). Biophilia: Does visual contact with nature impact on health and well-being? *International Journal of Environmental Research and Public Health, 6*(9), 2332–2343. <https://doi.org/10.3390/ijerph6092332>
- Heard, E. M., Mutch, A., Fitzgerald, L., & Pensalfini, R. (2013). Shakespeare in prison: Affecting health and wellbeing. *International Journal of Prisoner Health, 9*(3), 111–123. <https://doi.org/10.1108/IJPH-03-2013-0013>
- Harkrader, T., Burke, T. W., & Owen, S. S. (2004). Pound puppies: The rehabilitative uses of dogs in correctional facilities. *Corrections Today Magazine, 66*(2), 74–79.
- Hepper, E. G., Hart, C. M., Meek, R., Cisek, S., & Sedikides, C. (2014). Narcissism and empathy in young offenders and non-offenders. *European Journal of Personality, 28*(2), 201–210. <https://doi.org/10.1002/per.1939>
- Hughes, T., & Wilson, D. J. (2018). *Bureau of justice statistics: Reentry trends in the U.S.* <https://www.bjs.gov/content/reentry/reentry.cfm>

- Ingoglia, S., Coco, A. L., & Albiero, P. (2016). Development of a Brief form of the Interpersonal Reactivity Index (B-IRI). *Journal of Personality Assessment, 98*(5), 461–471. <https://doi.org/10.1080/00223891.2016.1149858>
- Jolliffe, D., & Farrington, D. P. (2004). Empathy and offending: A systematic review and meta-analysis. *Aggression and Violent Behavior, 9*(5), 441–476. <https://doi.org/10.1016/j.avb.2003.03.001>
- Kabacoff, R. I., Segal, D. L., Hersen, M., et al. (1997). Psychometric properties and diagnostic utility of the Beck Anxiety Inventory and the State-Trait Anxiety Inventory with older adult psychiatric outpatients. *Journal of Anxiety Disorders, 11*(1), 33–47. [https://doi.org/10.1016/S0887-6185\(96\)00033-3](https://doi.org/10.1016/S0887-6185(96)00033-3)
- Kaeble, D., & Cowhig, M. (2018). *Correctional populations in the United States, 2016*. Bureau of Justice Statistics. <https://www.bjs.gov/content/pub/pdf/cpus16.pdf>
- Lai, J. (1998). Pet facilitated therapy in correctional institutions. Office of the Deputy Commissioner for Women, Correctional Services of Canada. <http://www.cwhn.ca/en/node/25021>
- Lauterbach, O., & Hosser, D. (2007). Assessing empathy in prisoners—A shortened version of the Interpersonal Reactivity Index. *Swiss Journal of Psychology, 66*, 91–101. <https://doi.org/10.1024/1421-0185.66.2.91>
- Leonardi, R. J., Buchanan-Smith, H. M., McIvor, G., & Vick, S. J. (2017). “You think you’re helping them, but they’re helping you too”: Experiences of Scottish male young offenders participating in a dog training program. *International Journal of Environmental Research and Public Health, 14*(8), 945–972. <https://doi.org/10.3390/ijerph14080945>
- LeRoy, C. J., Bush, K., Trivett, J., & Gallagher, B. (2012). *The sustainability in prisons project: An overview (2004-2012)*. Gorham Publishing.
- Loinaz, I., Sanchez, L. M., & Vilella, A. (2018). Understanding empathy, self-esteem, and adult attachment in sexual offenders and partner-violent men. *Journal of Interpersonal Violence*. Advance online publication. <https://doi.org/10.1177/0886260518759977>
- Luszczynska, A., Gutiérrez-Doña, B., & Schwarzer, R. (2005). General self-efficacy in various domains of human functioning: Evidence from five countries. *International Journal of Psychology, 40*(2), 80–89. <https://doi.org/10.1080/00207590444000041>
- Luszczynska, A., Scholz, U., & Schwarzer, R. (2010). The General Self-Efficacy Scale: Multicultural validation studies. *The Journal of Psychology: Interdisciplinary and Applied, 139*(5), 439–457. <https://doi.org/10.3200/JRLP.139.5.439-457>
- Mackenzie, D. L. (1987). Age and adjustment to prison: Interactions with attitudes and anxiety. *Criminal Justice and Behavior, 14*(4), 427–447.
- Miller, P. A., & Eisenberg, N. (1988). The relation of empathy to aggressive and externalizing/antisocial behavior. *Psychological Bulletin, 103*(3), 324–344. <https://doi.org/10.1037/0033-2909.103.3.324>
- Minton, C. A., Perez, P. R., & Miller, K. (2015). Voices from behind prison walls: The impact of training service dogs on women in prison. *Society & Animals, 23*(5), 484–501. <https://doi.org/10.1163/15685306-12341379>

- Mulcahy, C., & McLaughlin, D. (2013). Is the tail wagging the dog? A Review of the Evidence for Prison Animal Programs. *Australian Psychologist*, *48*(5), 369–378. <https://doi.org/10.1111/ap.12021>
- Osborne, S. J., & Bair, R. (2003). Healing inmates' hearts and spirits with man's best friend. *Corrections Today*, *65*(2), 122–124.
- Perelman, A. M., Miller, S. L., Clements, C. B., Rodriguez, A., Allen, K., & Cavanaugh, R. (2012). Meditation in a deep south prison: A longitudinal study of the effects of Vipassana. *Journal of Offender Rehabilitation*, *51*(3), 176–198. <https://doi.org/10.1080/10509674.2011.632814>
- Richardson-Taylor, K., & Blanchette, K. (2001, September). *Results of an evaluation of the Pawsitive Directions Canine Program at Nova Institution for Women*. Ottawa, Ontario: Correctional Service of Canada. https://www.csc-scc.gc.ca/research/092/r108_e.pdf
- Schwarzer, R., & Jerusalem, M. (1995). Generalized Self-Efficacy scale. In J. Weinman, S. Wright, & M. Johnston (Eds.), *Measures in health psychology: A user's portfolio—Causal and control beliefs* (pp. 35–37). NFER-Nelson.
- Spielberger, C. D. (1972). *Anxiety: Current trends in research*. Academic Press.
- Spielberger, C. D. (1983). *Manual for the State-Trait Anxiety Inventory*. Consulting Psychologists Press.
- Spielberger, C. D., Gorsuch, R. L., & Lushene, R. E. (1970). *Manual for the State-Trait Anxiety Inventory*. Consulting Psychologists Press.
- Steiner, B., & Meade, B. (2016). Assessing the link between exposure to a violent prison context and inmate maladjustment. *Journal of Contemporary Criminal Justice*, *32*(4), 328–356. <https://doi.org/10.1177/1043986216660009>
- Strimple, E.O. (2003). A history of prison inmate-animal interaction programs. *American Behavioral Scientist*, *47*(1), 70–78. <https://doi.org/10.1177/0002764203255212>
- Van Langen, M., Wissink, I. B., Van Vugt, E. S., Van der Stouwe, T., & Stams, G. (2014). The relation between empathy and offending: A meta-analysis. *Aggression and Violent Behavior*, *19*(2), 179–189. <https://doi.org/10.1016/j.avb.2014.02.003>
- Van Wormer, J., Kigerl, A., & Hamilton, Z. (2017). Digging deeper: Exploring the value of prison-based dog handler programs. *The Prison Journal*, *97*(4), 520–538. <https://doi.org/10.1177/0032885517712481>
- Warren, J., Gelb, A., Horowitz, J., & Riordan, J. (2008, February). *One in 100: Behind bars in America 2008* (Pew Charitable Trust Report). https://www.pewtrusts.org/-/media/legacy/uploadedfiles/pcs_assets/2008/one20in20100pdf.pdf
- Washington State Department of Corrections. (2016). *Current programming*. <http://www.doc.wa.gov/corrections/programs/descriptions.htm>
- Washington State Institute for Public Policy. (2008). *Washington's offender accountability act: Department of Corrections' Static Risk Instrument*. http://www.wsipp.wa.gov/ReportFile/977/Wsipp_Washingtons-Offender-Accountability-Act-Department-of-Corrections-Static-Risk-Instrument_Full-Report-Updated-October-2008.pdf

- Wilson, E. O. (1984). Biophilia and the conservation ethic. In: S. Kellert S & E. O. Wilson (Eds.), *The biophilia hypothesis* (p. 57). Harvard University Press.
- Yogev, H. (2012). The development of empathy and group analysis. *Group Analysis*, 46(1), 61–80. <https://doi.org/10.1177/0533316412448312>

Author Biographies

Erin Flynn, MSW, is a research associate at the Institute for Human-Animal Connection, Graduate School of Social Work, University of Denver. Her research interests include corrections, quantitative methods, and positive development, with a focus on human–animal interactions (HAIs).

Katie Massey Combs, PhD, is a research associate with the Center for the Study and Prevention of Violence, University of Colorado Boulder. Her research focuses on the health and well-being of vulnerable young and on evaluation, prevention, and intervention programs.

Jaci Gandenberger, MSW, MAIS, is a research fellow at the Institute for Human-Animal Connection, Graduate School of Social Work, University of Denver. Her research interests lie at the intersections between human security, human rights, and animal and environmental well-being, particularly in cross-cultural contexts.

Philip Tedeschi, MSSW, is a clinical professor and executive director for The Institute for Human-Animal Connection, Graduate School of Social Work, University of Denver. His research and scholarship interests in the bio-affiliative connection between people and animals has spanned more than 27 years, focusing on many aspects of HAIs.

Kevin N. Morris, PhD, is a research associate professor in the Graduate School of Social Work, University of Denver. As the director of research for the University of Denver's Institute for Human-Animal Connection, he leads clinical research efforts to rigorously measure the impacts of animal-assisted interventions in environments such as family prevention services, residential treatment centers for adolescents, and prisons.